

KHOVANOV HOMOLOGY FOR ALTERNATING TANGLES

HERNANDO BURGOS SOTO

ABSTRACT. We describe a diagonal condition on the Khovanov complex of tangles, show that this condition is satisfied by the Khovanov complex of the single crossing tangles (\times) and (\times) , and prove that it is preserved by alternating planar algebra compositions. Hence, this condition is satisfied by the Khovanov complex of all alternating tangles. Finally, in the case of 0-tangles, that is links, our condition is equivalent to a well known result which states that the Khovanov homology of a non-split alternating link is supported in two lines.

CONTENTS

1. Introduction	1
2. Acknowledgement	3
3. The local Khovanov theory: Notation and some details	3
4. The category Kob_o and alternating planar algebras	10
5. Diagonal complexes	12
5.1. Applying unary operators	14
5.2. Applying binary operators	15
6. Proof of Theorem 1	17
7. Non-split alternating tangles and Lee's theorem	18
7.1. Proof of Theorem 2	18
References	20

1. INTRODUCTION

Khovanov [Kh] constructed an invariant of links which opened new prospects in knot theory and which is now known as the Khovanov homology. Bar-Natan in [BN] shows how to compute this invariant and found that it is a stronger invariant than the Jones polynomial. Khovanov, Bar-Natan and Garoufalidis [Ga] formulated several conjectures related to the Khovanov homology. One of these refers to the fact that the Khovanov homology of a non-split alternating link is supported in two lines. To see this, in Table 1, we present the dimension of the groups in the Khovanov homology for the Borromean link and illustrate that the no-zero dimension groups are located in two consecutive diagonals. The fact that every alternating link satisfies this property remained a conjecture until Lee proved it in [Lee].

Date: First edition: March. 22, 2010.

1991 Mathematics Subject Classification. 57M25.

Key words and phrases. Cobordism, Coherently diagonal complex Degree-shifted rotation number, De-looping, Gravity information, Khovanov homology, Diagonal complex, Planar algebra, Rotation number.

This work was partially supported by The Universidad del Norte in Barranquilla, Colombia and The Universidad Nacional in Bogota, Colombia.

$j \backslash i$	-3	-2	-1	0	1	2	3
7							1
5						2	
3						1	
1				4	2		
-1			2	4			
-3		1					
-5		2					
-7	1						

Table 1. The Khovanov homology for the Borromean link

In [BN1] Bar-Natan presented a new way of seeing the Khovanov homology. In his approach, a formal chain complex is assigned to every tangle. This formal chain complex, regarded within a special category, is an (up to homotopy) invariant of the tangle. For the particular case in which the tangle is a link, this chain complex coincides with the cube of smoothings presented in [Kh].

This local Khovanov theory was used in [BN2] to make an algorithm which provides a faster computation of the Khovanov homology of a link. The technique used in that last paper was also important for theoretical reasons. We can apply it to prove the invariance of the Khovanov homology, see [BN2]. It was also used in [BN3] to give a simple proof of Lee's result stated in [Lee1], about the dimension of the Lee variant of the Khovanov homology. Here, we will show how it can be used to state a generalization to tangles of the fore-mentioned Lee's theorem [Lee] about the Khovanov homology of alternating links.

We define a certain category \mathcal{Cob}_o^3 of *oriented cobordisms*. The objects of \mathcal{Cob}_o^3 are *oriented smoothing*, and the morphisms are oriented cobordisms. This orientation in the smoothings were previously utilized in [Bur] to define an integer parameter associated to the smoothing, and then generalize a Thistlethwaite result for the Jones polynomial stated in [Thi].

For an oriented smoothing σ , its *rotation number* is denoted by $R(\sigma)$. Specifically, for degree-shifted smoothings $\sigma\{q\}$ we define $R(\sigma\{q\}) := R(\sigma) + q$. We further use this *degree-shifted rotation number* to define a special class of chain complexes in $\text{Kom}(\text{Mat}(\mathcal{Cob}_o^3))$, of the form

$$\Omega : \quad \cdots \longrightarrow [\sigma_j^r]_j \longrightarrow [\sigma_j^{r+1}]_j \longrightarrow \cdots ,$$

which satisfies that for all degree-shifted smoothings $\sigma_j^r\{q\}$, $2r - R(\sigma_j^r\{q\})$ is a constant that we call *rotation constant* of the complex. In other words, twice the homological degrees and the degree-shifted rotation numbers of the smoothings always lie along a single diagonal. We call this type of chain complexes *diagonal complexes*. Furthermore, a *coherently diagonal* complex is a diagonal complex whose partial closure is also diagonal. Complexes of this type are the objects in the following theorem

Theorem 1. *Coherently diagonal complexes form an alternating planar algebra (that is, they are closed under “horizontal compositions” in alternating planar diagrams).*

Our second theorem follows from the first; for it reduces that proof to the simple task of verifying that the Khovanov homologies of the one-crossing tangles (\times) and (\times) (which are obviously alternating) are coherently diagonal:

Theorem 2. *Let T be a non-split alternating $2k$ -boundary tangle ($k > 0$), then the Khovanov homology $\text{Kh}(T)$ can be interpreted as a coherently diagonal complex.*

In the case of alternating tangles with no boundary, i.e., in the case of alternating links, this result reduces to Lee's theorem on the Khovanov homology of alternating links.

The work is organized as follows. In section 3, we review Bar-Natan local Khovanov theory and present two additional tools for the proof of theorem 1. These tools are propositions 3.5 and 3.7. Section 4 is devoted to introduce the category Cob_o^3 and give a quick review of some concepts related to alternating planar algebras. In particular we review the concepts of rotation number, alternating planar diagram, associated rotation number, and basic operators.

Section 5 introduces the concepts of diagonal complexes, coherently diagonal complexes, and their partial closures. We state here some results about the complexes obtained when a basic operator is applied to alternating elements, leading to the prove in section 6 of Theorem 1. Finally section 7 is dedicated to the study of non-split alternating tangles. Here, we prove Theorem 2 and derive from it Lee theorem formulated in [Lee].


2. ACKNOWLEDGEMENT

I wish to thank D. Bar-Natan, for many helpful conversations we had at the University of Toronto and for allowing me to use some figures from [BN1, BN2]. I would also like to thank N. Martin for their comments and suggestions.

3. THE LOCAL KHOVANOV THEORY: NOTATION AND SOME DETAILS

The notation and some results appearing here are treated in more details in [BN1, BN2, Naot]. Given a set B of $2k$ marked points on a circle C , a smoothing with boundary B is a union of strings a_1, \dots, a_n embedded in the plane disk for which C is the boundary, such that $\cup_{i=1}^n \partial a_i = B$. These strings are either closed curves, *loops*, or strings whose boundaries are points on B , *strands*. If $B = \emptyset$, the smoothing is a union of circles.

We denote $\text{Cob}^3(B)$, the category whose objects are smoothings with boundary B , and whose morphisms are cobordisms between such smoothings, regarded up to boundary preserving isotopy. The composition of morphisms is given by placing the second cobordism atop the other.



Our ground ring is one in which 2^{-1} exists. The dotted figure  is used as an abbreviation of $\frac{1}{2}$  and $\text{Cob}_{\bullet/l}^3(B)$ represents the category with the same objects and

morphisms as $Cob^3(B)$, whose morphisms are mod out by the local relations:

$$(1) \quad \begin{array}{c} \text{Diagram 1} = 0, \quad \text{Diagram 2} = 1, \quad \text{Diagram 3} = 0, \\ \text{and} \quad \text{Diagram 4} = \text{Diagram 5} \quad \text{Diagram 6} + \text{Diagram 7} = \text{Diagram 8}. \end{array}$$

We will use the notation \mathcal{Cob}^3 and $\mathcal{Cob}_{\bullet/l}^3$ as a generic reference, namely, $\mathcal{Cob}^3 = \bigcup_B Cob^3(B)$ and $\mathcal{Cob}_{\bullet/l}^3 = \bigcup_B Cob_{\bullet/l}^3(B)$. If B has $2k$ elements, we usually write $\mathcal{Cob}_{\bullet/l}^3(k)$ instead of $Cob_{\bullet/l}^3(B)$. If \mathcal{C} is any category, $\text{Mat}(\mathcal{C})$ will be the additive category whose object are column vectors (formal direct sums) whose elements are formal \mathbb{Z} -linear combinations of \mathcal{C} . Given two objects in this category,

$$\mathcal{O} = \begin{pmatrix} \mathcal{O}_1 \\ \mathcal{O}_2 \\ \vdots \\ \mathcal{O}_n \end{pmatrix} \quad \mathcal{O}^1 = \begin{pmatrix} \mathcal{O}_1^1 \\ \mathcal{O}_2^1 \\ \vdots \\ \mathcal{O}_m^1 \end{pmatrix},$$

the morphisms between these objects will be matrices whose entries are formal sums of morphisms between them. The morphisms in this additive category are added using the usual matrix addition and the morphism composition is modeled by matrix multiplication, i.e, given two appropriate morphisms $F = (F_{ik})$ and $G = (G_{kj})$ between objects of this category, then $F \circ G$ is given by

$$F \circ G = \sum_k F_{ik} G_{kj},$$

$\text{Kom}(\mathcal{C})$ will be the category of formal complexes over an additive category \mathcal{C} . $\text{Kom}_h(\mathcal{C})$ is $\text{Kom}(\mathcal{C})$ modulo homotopy. We also use the abbreviations $\text{Kob}(k)$ and $\text{Kob}_h(k)$ for denoting $\text{Kom}(\text{Mat}(\mathcal{Cob}_{\bullet/l}^3(k)))$ and $\text{Kom}_h(\text{Mat}(\mathcal{Cob}_{\bullet/l}^3(k)))$.

Objects and morphisms of the categories \mathcal{Cob}^3 , $\mathcal{Cob}_{\bullet/l}^3$, $\text{Mat}(\mathcal{Cob}_{\bullet/l}^3)$, $\text{Kob}(k)$, and $\text{Kob}_h(k)$ can be seen as examples of planar algebras, i.e., if D is a n -input planar diagram, it defines an operation among elements of the previously mentioned collections. See [BN1] for specifics of how D defines operations in each of these collections. In particular, if $(\Omega_i, d_i) \in \text{Kob}(k_i)$ are complexes, the complex $(\Omega, d) = D(\Omega_1, \dots, \Omega_n)$ is defined by

$$(2) \quad \begin{aligned} \Omega^r &:= \bigoplus_{r=r_1+\dots+r_n} D(\Omega_1^{r_1}, \dots, \Omega_n^{r_n}) \\ d|_{D(\Omega_1^{r_1}, \dots, \Omega_n^{r_n})} &:= \sum_{i=1}^n (-1)^{\sum_{j<i} r_j} D(I_{\Omega_1^{r_1}}, \dots, d_i, \dots, I_{\Omega_n^{r_n}}), \end{aligned}$$

$D(\Omega_1, \dots, \Omega_n)$ is used here as an abbreviation of $D((\Omega_1, d_1), \dots, (\Omega_n, d_n))$.

In [BN1] the following very desirable property is also proven. The Khovanov homology is a planar algebra morphism between the planar algebras $\mathcal{T}(s)$ of oriented tangles and $\text{Kob}_h(k)$. That is to say, for an n -input planar diagram D , and suitable tangles T_1, \dots, T_n , we have

$$(3) \quad Kh(D(T_1, \dots, T_n)) = D(Kh(T_1), \dots, Kh(T_n)).$$

This last property is used in [BN2] to show a local algorithm for computing the Khovanov

homology of a link. In that paper, Bar-Natan explained how it is possible to remove the loops in the smoothings, and some terms in the Khovanov complex $Kh(T_i)$ associated to the local tangles T_1, \dots, T_n , and then combine them together in an n -input planar diagram D obtaining $D(Kh(T_1), \dots, Kh(T_n))$, and the Khovanov homology of the original tangle.

The elimination of loops and terms can be done thanks to the following: Lemma 4.1 and Lemma 4.2 in [BN2]. We copy these lemmas verbatim:

Lemma 3.1. (*DeLooping*) *If an object S in $Cob_{\bullet/l}^3$ contains a closed loop ℓ , then it is isomorphic (in $Mat(Cob_{\bullet/l}^3)$) to the direct sum of two copies $S'\{+1\}$ and $S'\{-1\}$ of S in which ℓ is removed, one taken with a degree shift of $+1$ and one with a degree shift of -1 . Symbolically, this reads $\bigcirc \equiv \emptyset\{+1\} \oplus \emptyset\{-1\}$.*

The isomorphisms for the proof can be seen in:

$$(4) \quad \bigcirc \begin{array}{c} \nearrow \text{ (loop) } \\ \searrow \text{ (loop) } \end{array} \begin{array}{c} \left[\begin{array}{c} \bigcirc \{-1\} \\ \bigcirc \{+1\} \end{array} \right] \begin{array}{c} \nearrow \text{ (loop) } \\ \searrow \text{ (loop) } \end{array} \bigcirc$$

using all the relations in (1).

Lemma 3.2. (*Gaussian elimination, made abstract*) *If $\phi : b_1 \rightarrow b_2$ is an isomorphism (in some additive category \mathcal{C}), then the four term complex segment in $Mat(\mathcal{C})$*

$$(5) \quad \dots [C] \xrightarrow{\begin{pmatrix} \alpha \\ \beta \end{pmatrix}} \begin{bmatrix} b_1 \\ D \end{bmatrix} \xrightarrow{\begin{pmatrix} \phi & \delta \\ \gamma & \epsilon \end{pmatrix}} \begin{bmatrix} b_2 \\ E \end{bmatrix} \xrightarrow{\begin{pmatrix} \mu & \nu \end{pmatrix}} [F] \dots$$

is isomorphic to the (direct sum) complex segment

$$(6) \quad \dots [C] \xrightarrow{\begin{pmatrix} 0 \\ \beta \end{pmatrix}} \begin{bmatrix} b_1 \\ D \end{bmatrix} \xrightarrow{\begin{pmatrix} \phi & 0 \\ 0 & \epsilon - \gamma\phi^{-1}\delta \end{pmatrix}} \begin{bmatrix} b_2 \\ E \end{bmatrix} \xrightarrow{\begin{pmatrix} 0 & \nu \end{pmatrix}} [F] \dots$$

Both these complexes are homotopy equivalent to the (simpler) complex segment

$$(7) \quad \dots [C] \xrightarrow{(\beta)} [D] \xrightarrow{(\epsilon - \gamma\phi^{-1}\delta)} [E] \xrightarrow{(\nu)} [F] \dots$$

Here C , D , E and F are arbitrary columns of objects in \mathcal{C} and all Greek letters (other than ϕ) represent arbitrary matrices of morphisms in \mathcal{C} (having the appropriate dimensions, domains and ranges); all matrices appearing in these complexes are block-matrices with blocks as specified. b_1 and b_2 are billed here as individual objects of \mathcal{C} , but they can equally well be taken to be columns of objects provided (the morphism matrix) ϕ remains invertible.

It will be useful for our purpose to enunciate also the following lemma which is easily demonstrable using the obvious morphism of complexes.

Lemma 3.3. *If B is an object of $Mat(\mathcal{C})$ involved in a chain complex Ω , then it is possible to interchange the position of two elements b_i, b_j of B obtaining a homotopy equivalent complex. This interchange also changes the position of the i -th and j -th rows of the morphism pointing*

at B and the i -th and j -th columns of the morphism coming from B . In other words, the three term complex segment in $\text{Mat}(\mathcal{C})$

$$(8) \quad \cdots \begin{bmatrix} a_1 \\ \vdots \\ a_n \end{bmatrix} \xrightarrow{\begin{pmatrix} \alpha_{11} & \cdots & \alpha_{1n} \\ \vdots & & \vdots \\ \alpha_{i1} & \cdots & \alpha_{in} \\ \vdots & & \vdots \\ \alpha_{j1} & \cdots & \alpha_{jn} \\ \vdots & & \vdots \\ \alpha_{m1} & \cdots & \alpha_{mn} \end{pmatrix}} \begin{bmatrix} b_1 \\ \vdots \\ b_i \\ \vdots \\ b_j \\ \vdots \\ b_m \end{bmatrix} \xrightarrow{\begin{pmatrix} \beta_{11} & \cdots & \beta_{1i} & \cdots & \beta_{1j} & \cdots & \beta_{1m} \\ \vdots & & \vdots & & \vdots & & \vdots \\ \beta_{p1} & \cdots & \beta_{pi} & \cdots & \beta_{pj} & \cdots & \beta_{pm} \end{pmatrix}} \begin{bmatrix} c_1 \\ \vdots \\ c_p \end{bmatrix} \cdots$$

is isomorphic to

$$(9) \quad \cdots \begin{bmatrix} a_1 \\ \vdots \\ a_n \end{bmatrix} \xrightarrow{\begin{pmatrix} \alpha_{11} & \cdots & \alpha_{1n} \\ \vdots & & \vdots \\ \alpha_{j1} & \cdots & \alpha_{jn} \\ \vdots & & \vdots \\ \alpha_{i1} & \cdots & \alpha_{in} \\ \vdots & & \vdots \\ \alpha_{m1} & \cdots & \alpha_{mn} \end{pmatrix}} \begin{bmatrix} b_1 \\ \vdots \\ b_j \\ \vdots \\ b_i \\ \vdots \\ b_m \end{bmatrix} \xrightarrow{\begin{pmatrix} \beta_{11} & \cdots & \beta_{1j} & \cdots & \beta_{1i} & \cdots & \beta_{1m} \\ \vdots & & \vdots & & \vdots & & \vdots \\ \beta_{p1} & \cdots & \beta_{pj} & \cdots & \beta_{pi} & \cdots & \beta_{pm} \end{pmatrix}} \begin{bmatrix} c_1 \\ \vdots \\ c_p \end{bmatrix} \cdots$$

where every Latin and Greek letter represents respectively a smoothing or a cobordism.

From the three previous lemmas we infer that the Khovanov complex of a tangle is homotopy equivalent to a chain of complex without loops in the smoothings, and in which every differential is a non-invertible cobordism. In other words, if (Ω, d) is a complex in $\mathcal{Cob}_{\bullet/l}^3$, we can use lemmas 3.1, 3.2 and 3.3, and obtain a homotopy equivalent chain complex (Ω', d') with no loop in its smoothings and no invertible cobordism in its differentials. We say that (Ω', d') is a *reduced complex* of (Ω, d) .

For our purposes, it will be useful to recall here the concept of *bounded* chain complex. See [Wei]. A chain complex

$$\Omega : \quad \cdots \Omega^r \xrightarrow{d^r} \Omega^{r+1} \cdots$$

is called *bounded* if almost all the Ω^r are zero. If $\Omega^{r_0} \neq 0$, $\Omega^{r_M} \neq 0$ and $\Omega^r = 0$ unless $r_0 \leq r \leq r_M$, we say that (Ω, d) has *amplitude* in $[r_0, r_M]$.

Definition 3.4. Let (Ω, d) be a bounded chain complex in Kob with amplitude in $[r_0, r_M]$.

Let $\Omega^r = \begin{bmatrix} \sigma_1^r \\ \vdots \\ \sigma_{n_r}^r \end{bmatrix}$ be the vector in the complex (Ω, d) with homological degree r . Thus the set S^r formed by the elements of this vector has cardinal n_r . Assume that the cardinal of $S = \bigcup_{r=r_0}^{r_M} S^r$ is N , that is to say, there are in total N smoothings in the complex. A *numeration* of (Ω, d) is a map $g : S \rightarrow \{1, \dots, N\}$ defined in this way: $g(\sigma_1^{r_0}) = 1$; $g(\sigma_1^{r+1}) = g(\sigma_{n_r}^r) + 1$, if $r_0 \leq r < r_M$; and $g(\sigma_{i+1}^r) = g(\sigma_i^r) + 1$, if $1 \leq i < n_t$. This numerates the smoothings in (Ω, d) , and we can rewrite σ_i^r as $\sigma_{g(i)}^r$.

Given a complex (Ω, d) then the component of d connecting σ_j and σ_i is denoted d_{ij} . It is clear from the definition of a numeration in (Ω, d) that if $i \leq j$, then $d_{ij} : \sigma_j \rightarrow \sigma_i$ is the zero cobordism. The Figure 1 displays an example of a complex with its numeration.




Figure 1. A numeration in a complex, the dotted circles around the smoothings represent the discs in which the smoothings are embedded. The subindex in each smoothing is the number assigned to this smoothing by the numeration.

Proposition 3.5. Let (Ψ, e) and (Φ, f) be chain complexes in Kob, Let (Φ, f) be a bounded complex in $[t_0, t_M]$, and D an appropriate 2-input planar arc diagram. Let ϕ_1, \dots, ϕ_N be a numeration of (Φ, f) Then $D(\Psi, \Phi)$ is homotopy equivalent to a chain complex (Ω, d) with the following properties:

- (1) Every vector Ω^r is of the form

$$\Omega^r = \bigoplus_{\substack{t_0 \leq t \leq t_M \\ s = r - t}} D(\Psi^s, \Phi^t)$$

can be regarded as a block column matrix $\begin{pmatrix} \Omega_1^r \\ \vdots \\ \Omega_N^r \end{pmatrix}$ in which each block $\Omega_i^r = D(\Psi^s, \phi_i)$,

where ϕ_i is a smoothing in Φ^t .

- (2) The differential matrixes d^r can be seen as lower block triangular matrices with blocks $d_{ij}^r : \Omega_j^r \rightarrow \Omega_i^{r+1}$.

Proof. The first of these statements follows immediately from the definition of $D(\Psi, \Phi)$, equations (2). Obviously, if $s \leq s_0$ or $s \geq s_M$ we consider $\Psi^s = 0$.

For the second statement we see that given $r = s + t$, the matrix d^r is defined by the second of the equations (2), and is given by

$$(10) \quad d|_{D(\Psi^s, \Phi^t)} = D(e, I_{\Phi^t}) + (-1)^s D(I_{\Psi^s}, f).$$


This matrix can be seen as a block matrix in which each block d_{ij}^r is a morphism of the form $d_{ii}^r : D(\Psi^s, \phi_i) \rightarrow D(\Psi^{s+1}, \phi_i)$, and any other block is a morphism of the form $d_{ij}^r : D(\Psi^s, \phi_j) \rightarrow D(\Psi^{s+1}, \phi_i)$ with $i \neq j$. We conclude from this that $D(e, I_{\Phi^t})$ in the right side of equation (10) is concentrated in the diagonal of blocks. It is clear that the blocks over the diagonal are zero, since they are part of $\pm D(I_{\Psi^s}, f)$ in the right side of equation (10) and if $i < j$, $f_{ij} : \phi_j \rightarrow \phi_i$ is the zero cobordism. \square

Remark 3.6. The blocks $\Omega_i^r = D(\Psi^s, \phi_i)$ in Ω^r , and the blocks $d_{ii}^r : D(\Psi^s, \phi_i) \rightarrow D(\Psi^{s+1}, \phi_i)$ in the diagonal of d^r (here $s = r - t$ and ϕ_i is a smoothing in Φ^t), determine the complex

$$D(\Psi, \phi_i) = \cdots \Omega_{ii}^r \xrightarrow{d^r} \Omega_{ii}^{r+1} \cdots$$

\square

We illustrate the previous proposition with an example. Let D be the binary operator defined from the planar arc diagram of the right. If we place the complex



$$\Psi = \left[\begin{array}{c} \text{Diagram 1} \\ \text{Diagram 2} \end{array} \right] \xrightarrow{\left[\begin{array}{c} \text{Diagram 3} \\ \text{Diagram 4} \end{array} \right]} \left[\begin{array}{c} \text{Diagram 5} \\ \text{Diagram 6} \end{array} \right] \xrightarrow{\left[\begin{array}{c} \text{Diagram 7} \\ \text{Diagram 8} \end{array} \right]} \left[\begin{array}{c} \text{Diagram 9} \end{array} \right]$$

— -1 ————— 0 ————— 1 —

in the first entry of D and

$$\Phi = \left[\begin{array}{c} \text{Diagram 10} \\ \text{Diagram 11} \end{array} \right] \xrightarrow{\left[\begin{array}{c} \text{Diagram 12} \end{array} \right]} \left[\begin{array}{c} \text{Diagram 13} \end{array} \right]$$

— 0 ————— 1 —

$$D(\Psi, \Phi) = \left[\begin{array}{c} \text{Diagram 1} \\ 0 \end{array} \right] \xrightarrow{d^{-1}} \left[\begin{array}{c} \text{Diagram 2} \\ \text{Diagram 3} \\ \text{Diagram 4} \end{array} \right] \xrightarrow{d^0} \left[\begin{array}{c} \text{Diagram 5} \\ \text{Diagram 6} \\ \text{Diagram 7} \end{array} \right] \xrightarrow{d^1} \left[\begin{array}{c} 0 \\ \text{Diagram 8} \end{array} \right]$$

$\text{--- } -1 \text{ --- } 0 \text{ --- } 1 \text{ --- } 2 \text{ ---}$

$$\begin{aligned}
d^{-1} &= \begin{bmatrix} \left[\begin{array}{c} \text{diagram 1} \end{array} \right] \begin{bmatrix} 0 \end{bmatrix} \\ \left[\begin{array}{c} \text{diagram 2} \end{array} \right] \begin{bmatrix} 0 \end{bmatrix} \\ \left[\begin{array}{c} \text{diagram 3} \end{array} \right] \begin{bmatrix} 0 \end{bmatrix} \end{bmatrix} & \quad d^0 = \begin{bmatrix} \left[\begin{array}{cc} \text{diagram 4} & \text{diagram 5} \end{array} \right] \begin{bmatrix} 0 \end{bmatrix} \\ \left[\begin{array}{cc} \text{diagram 6} & 0 \end{array} \right] \begin{bmatrix} \text{diagram 7} \end{bmatrix} \\ \left[\begin{array}{cc} 0 & \text{diagram 8} \end{array} \right] \begin{bmatrix} \text{diagram 9} \end{bmatrix} \end{bmatrix} \\
d^1 &= \begin{bmatrix} \left[\begin{array}{cc} 0 & 0 \end{array} \right] \\ \left[\begin{array}{cc} \text{diagram 10} & \text{diagram 11} \end{array} \right] \end{bmatrix}
\end{aligned}$$

Proposition 3.7. *Assume that the three differential matrices in the four term complex segment (5) of lemma 3.2 are block-lower-triangular matrices. After applying gauss elimination, the resulting three differential matrices in the four term complex segment (7) are also block-lower-triangular matrices. Furthermore, the lowest right block of the three initial differential matrices remained unchanged after the Gauss elimination.*

Proof. It is clear that if $\begin{pmatrix} \alpha \\ \beta \end{pmatrix}$, $\begin{pmatrix} \phi & \delta \\ \gamma & \epsilon \end{pmatrix}$, and $\begin{pmatrix} \mu & \nu \end{pmatrix}$ are block-lower-triangular matrices, so they are β , ϵ , and ν . Therefore, it is clear that after Gauss elimination, the first and the third of the differential matrices in the form term complex (7) are block-lower-triangular matrices with the same initial lowest-right block.

To prove that the same happens with the second block, we observe that if $\begin{pmatrix} \phi & \delta \\ \gamma & \epsilon \end{pmatrix}$ is a block-lower-triangular matrices then $\begin{pmatrix} \phi & \delta \\ \gamma & \epsilon \end{pmatrix} = \begin{pmatrix} \phi & \delta_1 & 0 \\ \gamma_1 & \epsilon_1 & 0 \\ \gamma_2 & \epsilon_2 & \epsilon_3 \end{pmatrix}$; where $\delta = \begin{pmatrix} \delta_1 & 0 \end{pmatrix}$, $\gamma = \begin{pmatrix} \gamma_1 \\ \gamma_2 \end{pmatrix}$, and $\epsilon = \begin{pmatrix} \epsilon_1 & 0 \\ \epsilon_2 & \epsilon_3 \end{pmatrix}$. Each 0 in the previous matrices is actually a block of zeros.

An immediate consequence of the previous paragraph is that the second differential matrix in the four term complex segment (7) is given by


$$\epsilon - \gamma\phi^{-1}\delta = \begin{pmatrix} \epsilon_1 - \gamma_1\phi^{-1}\delta_1 & 0 \\ \epsilon_2 - \gamma_2\phi^{-1}\delta_1 & \epsilon_3 \end{pmatrix}.$$

This completes the proof. \square

4. THE CATEGORY Kob_o AND ALTERNATING PLANAR ALGEBRAS


We introduce an alternating orientation in the objects of $\text{Cob}_{\bullet/l}^3(k)$. This orientation induces an orientation in the cobordisms of this category. These oriented k -strand smoothings and cobordisms form the objects and morphisms in a new category. The composition between cobordisms in this oriented category is defined in the standard way, and it is regarded as a graded category, in the sense of [BN1, Section 6]. We subject out the cobordisms in this oriented category to the relations in (1) and denote it as $\text{Cob}_o^3(k)$. Now we can follow [BN1] and define sequentially the categories, $\text{Mat}(\text{Cob}_o^3(k))$, $\text{Kom}(\text{Mat}(\text{Cob}_o^3(k)))$ and $\text{Kom}_{/h}(\text{Mat}(\text{Cob}_o^3(k)))$. This last two categories are what we denote $\text{Kob}_o(k)$, and $\text{Kob}_{o/h}$. As usual, we use Kob_o , and $\text{Kob}_{o/h}$, to denote $\bigcup_k \text{Kob}_o(k)$ and $\bigcup_k \text{Kob}_{o/h}(k)$ respectively.

We denote the class of oriented smoothings as \mathcal{S}_o . An alternatively oriented d -input planar diagram, see [Bur], provides a good tool for the horizontal composition of objects in \mathcal{S}_o , Cob_o^3 , $\text{Mat}(\text{Cob}_o^3(k))$, Kob_o , and $\text{Kob}_{o/h}$. The orientation in the diagrams can be provided as in the figure at the right. For making this text a little more self-contained, we are going to recall briefly some concepts presented before in [Bur]. Given oriented smoothings $\sigma_1, \dots, \sigma_d$, a suitable alternating d -input planar diagram D to compose them has the property that the i -th input disc has as many boundary points as σ_i . Moreover placing σ_i in the i -th input disc, The orientation (the coloring) of σ_i and D match.



Given an open strand α of an alternating oriented smoothing σ , possibly with loops, enumerate the boundary points of σ in such a way that α can be denoted by $(0, i)$. The *rotation number* of $(0, i)$, $R(\alpha)$, is $\frac{i-k}{2k}$. If α is a loop, $R(\alpha) = 1$ if α is oriented counterclockwise, and

$R(\alpha) = -1$ if α is oriented clockwise. The rotation number of σ is the sum of the rotation numbers of its strings. See figure 3 We are going to use this alternating diagrams to compute




$$R(\alpha) = \frac{1-3}{6}$$

Figure 3. $\alpha = (0, 1)$, $R(\alpha) = -\frac{2}{6} = -\frac{1}{3}$. The rotation number of the complete resolution is 0

non-split alternating tangles, and we want to preserve the non-split property of the tangle. Hence, it will be better if we use d -input type \mathcal{A} diagrams.

A d -input type \mathcal{A} diagram has an even number of strings ending in each of its boundary components, and every string that begins in the external boundary ends in a boundary of an internal disk. We can classify the strings as: *curls*, if they have its ends in the same input disc; *interconnecting arcs*, if its ends are in different input discs, and *boundary arcs*, if they have one end in an input disc and the other in the external boundary of the output disc. The arcs and the boundaries of the discs divide the surface of the diagram into disjoint regions. Some arcs and regions will be useful in the following definitions and propositions.



Definition 4.1. We assign the following numbers to every d -input planar diagram D :

- i_D : number of interconnecting arcs and curls, i.e., the number of non-boundary arcs.
- w_D : number of negative internal regions. That is, in the checkerboard coloring, the white regions whose boundary does not meet the external boundary of D .
- R_D : the rotation associated number, which is given by the formula

$$R_D = \frac{1}{2}(1 + i_D - d) - w_D$$

Proposition 4.2. Given the smoothings $\sigma_1, \dots, \sigma_d$ and a suitable d -input planar diagram D , where every smoothing can be placed, the rotation number of $D(\sigma_1, \dots, \sigma_d)$ is:

$$(11) \quad R(D(\sigma_1, \dots, \sigma_d)) = R_D + \sum_{i=1}^d R(\sigma_i)$$

Definition 4.3. An alternating planar algebra is a triplet $\{\mathcal{P}, \mathcal{D}, \mathcal{O}\}$ in which \mathcal{P} , \mathcal{D} , and \mathcal{O} have the same properties as in the definition of a planar algebra but with the collection \mathcal{D} containing only \mathcal{A} -type planar diagrams.

Diagrams with only one or two input discs deserves special attention. Operators defined from diagram like these are very important for our purposes since some of them are considered as the generators of the entire collection of operators in a connected alternating planar algebra.




Figure 4. Examples of basic planar diagrams

Definition 4.4. A basic planar diagram is a 1-input alternating planar diagram with a curl in it, or a 2-input alternating planar diagram with only one interconnecting arc. A basic operator is one defined from a basic planar diagram. A negative unary basic operator is one defined from a basic 1-input diagram where the curl completes a negative loop. A positive unary basic operator is one defined from a basic 1-input diagram where the curl completes a positive loop. A binary operator is one defined from a basic 2-input planar diagram.

Proposition 4.5. *The rotation associated number of a planar diagram belongs to $\frac{1}{2}\mathbb{Z}$ and the case when we have a basic planar diagram it is given as follows:*

- If D is a negative unary basic operator, $R_D = -\frac{1}{2}$
- If D is a binary basic operator, $R_D = 0$
- If D is a positive unary basic operator, $R_D = \frac{1}{2}$

Proposition 4.6. *Any operator D in an alternatively oriented planar algebra is the finite composition of basic operators.*

5. DIAGONAL COMPLEXES

Once we have applied lemma 3.1 to an element of Kob_o , we obtain a complex (Ω, d) , which preserves some properties of the former one, but with a change in the rotation number of the element $\sigma\{q_\sigma\}$, in which we have applied the delooping. In fact, the smoothing has been replaced in the complex by a couple whose rotation number has changed either by -1 or by +1. This shift in the rotation number could be even greater if we continue removing loops in the same smoothing. From lemma 3.1, we know that there is also a change in the grading shift of the smoothings. So it would be a good idea to define a concept that states a relation between the rotation number of σ and its grading shift q_σ .

Definition 5.1. Let (Ω, d) be a class-representative of $\text{Kob}_{o/h}$, and let $\sigma_i\{q_i\}$ be a shifted degree object in Ω^r , then its degree-shifted rotation number is $\overline{R}(\sigma_i\{q_i\}) = R(\sigma_i) + q_i$

Definition 5.2. A diagonal complex is a degree-preserving differential chain complex (Ω, d)

$$\dots \Omega^r \xrightarrow{d^r} \Omega^{r+1} \dots$$

in Kob_o , satisfying that for each homological degree r and each shifted degree object $\sigma_i\{q_i\}$ in Ω^r , we have that $2r - \overline{R}(\sigma_i\{q_i\}) = C_\Omega$, where C_Ω is a constant that we call rotation constant of (Ω, d) .

Here we have some examples of diagonal complexes in Kob_o .

Example 5.3. As in [BN2], a dotted line represent a dotted curtain, and \succ stands for the saddle $\succ \longrightarrow \prec$

(1)

$$\Omega_1 = \left(\begin{array}{c} \text{dotted curtain} \\ \text{saddle} \end{array} \right) \{-2\} \xrightarrow{\succ} \left(\begin{array}{c} \text{dotted curtain} \\ \text{saddle} \end{array} \right) \{-1\} .$$

This is the Khovanov homology of the negative crossing \succ , now with orientation in the smoothings. Remember that the first term has homological degree -1. In this example the rotation number in the first term is $-\frac{1}{2}$ and in the second term it is $\frac{1}{2}$. Observe that in each case, the difference between 2 times the homological degree r and the shifted rotation number is $\frac{1}{2}$.




Figure 5. A diagonal complex.

- (2) In Figure 5, the number below each smoothing is the grading shift of the smoothing. The upper line below the complex represents the homological degree r , and the lower one represents the degree-shifted rotation number. For instance, the rotation number in the first smoothing with homological degree 1 has rotation number 0 and a grading shift by -1. In the second smoothing of the same vector, the rotation number is -1 and its grading shift is 0, so both term has the same degree-shifted rotation number. We see in this example, that for each r we have that $2r - R = 3$, so this is a diagonal complex.


Now, we can establish a parallel between what we did with alternating elements in $\mathcal{M}_k^{(o)}$ and diagonal complexes in Kob_o in such a way that we can obtain similar results as those obtained in section 4 of [Bur].

5.1. Applying unary operators. The reduced complexes in $\text{Kom}(\text{Mat}(\text{Cob}_o^3))$ can be inserted in appropriate unary basic planar diagrams, and then apply lemmas 3.1, 3.2, and 3.3 to obtain again a reduced complex in Kob_o . This process can be summarized in the following steps:

- (1) placing of the complex in the corresponding input disc of the d -input planar arc diagram by using equations (2),
- (2) removing the loops obtained by applying lemma 3.1, i.e, replacing each of them by a copy of $\emptyset\{+1\} \oplus \emptyset\{-1\}$, and
- (3) applying lemma 3.3 and gaussian elimination (lemma 3.2), and removing in this way each invertible differential in the complex.

Definition 5.4. Let (Ω, d) be a chain complex in $\text{Kom}(\text{Mat}(\text{Cob}_o^3(k)))$, then a partial closure of (Ω, d) is a chain complex of the form $D_l \circ \cdots \circ D_1(\Omega)$ where $0 \leq l < k$ and every D_i ($1 \leq i \leq l$) is a unary basic operator

We have diagonal complexes whose partial closures are again diagonal complexes. For instance, embedding Ω_1 of the example 5.3 in a unary basic planar diagram U_1 as the one on the right which has an associated rotation number $R_{U_1} = \frac{1}{2}$, produces the chain complex.



$$\begin{aligned}
 U_1(\Omega_1) &= \left[\text{diagram of two loops with dots and arrows} \right] \{-2\} \xrightarrow{[\boxtimes]} \left[\text{diagram of two loops with dots and arrows} \right] \{-1\} \\
 &\sim \left[\text{diagram of two loops with dots and arrows} \right] \{-2\} \xrightarrow{\left[\begin{smallmatrix} \text{diagram of a loop with a dot and arrow} \\ \text{diagram of a loop with a dot and arrow} \end{smallmatrix} \right]} \left[\begin{smallmatrix} \text{diagram of a loop with a dot and arrow} \{-2\} \\ \text{diagram of a loop with a dot and arrow} \{0\} \end{smallmatrix} \right]
 \end{aligned}$$

The last complex is the result of applying lemma 3.1. Applying now lemma 3.2 we obtain a homotopy equivalent complex

$$U_1(\Omega_1) \sim 0 \xrightarrow{[0]} \left[\text{diagram of a loop with a dot and arrow} \right] \{0\}$$

which is also a diagonal complex, but now with rotation constant zero.


Definition 5.5. Let (Ω, d) be a bounded diagonal complex in Kob_o with rotation constant C_R . We say that (Ω, d) is *coherently diagonal* if for any appropriated unary operator with associated rotation number R_U , the closure $U(\Omega, d)$ has a reduced form which is a diagonal complex with rotation constant $C_R - R_U$.

We denote as $\mathcal{D}(k)$ the collection of all coherently diagonal complexes in $\text{Kom}(\text{Mat}(\text{Cob}_o^3(k)))$, and as usual, we write \mathcal{D} to denote $\bigcup_k \mathcal{D}(k)$. It is easy to prove that any coherently diagonal complex satisfies that:

- (1) after delooping any of the positive loops obtained in any of its partial closure, by using lemma 3.2, the negative shifted-degree term can be eliminated.

- (2) after delooping any of the negative loops obtained in any of its partial closure, by using lemma 3.2, the positive shifted-degree term can be eliminated.

Since the computation of any other of its partial closures produces other diagonal complex, the complex Ω_1 of the example 5.3 is an element of $\mathcal{D}(2)$. Another example of coherently diagonal complex is the complex Ω_2 of the same example. This last complex has $C_R = 3$. All of its partial closures $U(\Omega_2)$ are diagonal complexes with rotation constant given by $C_R - R_U$. Here, we only calculate the one produced by inserting the element in the closure disc U , with $R_U = -\frac{1}{2}$, that appears on the right.



It will be easy for the reader to compute the other partial closures. Inserting Ω_2 in U produces the complex of Figure 6, which is also a diagonal complex, but with a loop in some




Figure 6. A diagonal complex inserted in a negative unary basic diagram U .

of its smoothings. Observe that the rotation number of the smoothings have decreased in $\frac{1}{2}$ after having been inserted in a negative unary basic diagram.

After applying lemmas 3.1 and 3.2, we obtain the complex in Figure 7 which is also a diagonal complex, but now with rotation constant $\frac{7}{2}$.

5.2. Applying binary operators.

Proposition 5.6. *If D is an appropriate binary basic operator and $(\Psi, e), (\Phi, f)$ are diagonal complexes in Kob_o with rotation constants C_Ψ and C_Φ respectively, then $D(\Psi, \Phi)$ is a diagonal complex with rotation constant $C_\Psi + C_\Phi$.*




Figure 7. A partial closure of a coherently diagonal complex is also a diagonal complex.

Proof. Inserting (Ψ, e) and (Φ, f) in the disc D produces the complex $(\Omega, d) = D(\Psi, \Phi)$, which by equation (2) satisfies

$$(12) \quad \Omega^r = \bigoplus_{r=s+t} D(\Psi^s, \Phi^t)$$

and

$$(13) \quad d|_{D(\Psi^s, \Phi^t)} = D(e, I_{\Phi^t}) + (-1)^s D(I_{\Psi^s}, f)$$

If $\psi\{q_\psi\}$ and $\phi\{q_\phi\}$ are respectively elements in the vectors Ψ^s and Φ^t , so by equation (12) the elements in the vector Ω^r are of the form $D(\psi, \phi)\{q_\psi + q_\phi\}$. As ψ and ϕ are smoothings with no loops, the same we have for $D(\psi, \phi)$ and by using propositions 3.7 and 3.10 in [Bur], we obtain

$$R(D(\psi, \phi)) + q_\psi + q_\phi = R(\psi) + R(\phi) + q_\psi + q_\phi.$$

Therefore, the homological degree r is given by $s - (R(\psi) + q_\psi) + t - (R(\phi) + q_\phi) = C_\Psi + C_\Phi$ \square

Proposition 5.7. *Let (Ψ, e) and (Φ, e) complex in \mathcal{D} with rotation constant C_Ψ and C_Φ respectively, and let D be a binary basic planar operator in which $D(\Psi, \Phi)$ is well defined. For each partial closure $C(D(\Psi, \Phi))$, there exists an operator D' defined on a diagram without curls and chain complexes Ψ', Φ' in \mathcal{D} such that*

$$C(D(\Psi, \Phi)) = D'(\Psi', \Phi')$$

Proof. The proof similar to the proof of proposition 4.7 in [Bur] \square

Proposition 5.8. *Let σ and τ be smoothings, and let D be a suitable binary planar operator defined from a no-curl planar arc diagram with output disc D_0 , input discs D_1, D_2 , associated*

rotation constant R_D and with at least one boundary arc ending in D_1 , then there exists a closure operator C and a unary operator D' defined from a no-curl planar arc diagram such that $D(\sigma, \tau) = D'(C(\sigma))$. Moreover, if $(\Omega, d) \in \mathcal{D}$ has rotation constant C_Ω , then $D(\Omega, \tau)$ is a diagonal complex with rotation constant $C_\Omega - R(\tau) - R_D$.

Proof. To prove that there exists a closure operator C and a unary operator D' defined from a no-curl planar arc diagram such that $D(\sigma, \tau) = D'(C(\sigma))$, we reason as in the proof of proposition 4.8 in [Bur]. To prove that the rotation constant of $D(\Omega, \tau)$ is $C_\Omega - R(\tau) - R_D$, we observe that for each smoothing $\sigma\{q_\sigma\}$ in Ω the shifted rotation number satisfies $\overline{R}(D(\sigma\{q_\sigma\} \tau)) = R_D + \overline{R}(\sigma\{q_\sigma\}) + R(\tau) = R_D + 2r - C_\Omega + R(\tau)$. Therefore, $2r - \overline{R}(D(\sigma\{q_\sigma\} \tau)) = C_\Omega - R(\tau) - R_D$ \square

6. PROOF OF THEOREM 1

Before proving the first main theorem, let us state the following result.

Lemma 6.1. *Let (Ψ, e) and (Φ, f) be a coherently diagonal complexes with rotation constants C_Ψ and C_Φ respectively. Suppose that D is a binary operator with no curls and with associated rotation constant R_D , then $D(\Psi, \Phi)$ is a diagonal complex with rotation constant $C_\Psi + C_\Phi - R_D$.*

Proof. Assume that (Φ, f) is bounded in $[t_0, t_M]$. Let $g : S \rightarrow \{1, \dots, N\}$ a numeration in (Φ, f) , we apply induction on N . For the case $N=1$, the result is obvious by proposition 5.8.

Assume that the statement is valid for any diagonal complex with numeration $g : S \rightarrow \{1, \dots, N-1\}$. $(\Omega, d) = D(\Psi, \Phi)$, let (Φ', f') be the complex resulting from eliminating in (Φ, f) , the last smoothing ϕ_N and every cobordism that have ϕ_N as the image. It will be easy for the reader to prove that (Ψ', f') is in fact a chain complex. By proposition 3.5, we have that the complex $D(\Psi, \Phi)$ is formed by segments of the form

$$(14) \quad \dots [\Omega^{r-1}] \xrightarrow{\begin{pmatrix} d^{r-1} & 0 \\ \rho^{r-1} & d_n^{s-1} \end{pmatrix}} \begin{bmatrix} \Omega'^r \\ \Omega_n^s \end{bmatrix} \xrightarrow{\begin{pmatrix} d^r & 0 \\ \rho^r & d_n^s \end{pmatrix}} \begin{bmatrix} \Omega'^{r+1} \\ \Omega_n^{s+1} \end{bmatrix} \xrightarrow{\begin{pmatrix} d^{r+1} & 0 \\ \rho^{r+1} & d_n^{s+1} \end{pmatrix}} [\Omega^{r+2}] \dots$$

Here, $(\Omega', d') = D(\Psi, \Phi')$, $d_n^s = D(e^s, I_{\Phi_n})$, $s = r - t_M$, and $\rho^r = (-1)^s D(I_{\Psi^s}, f_n)$, where f_n is the component of f that has ϕ_n as the image.

By the induction hypothesis, $D(\Phi, \Psi')$ is a coherently alternating complex, so it is possible to carry out delooping and gauss eliminations in $D(\Phi, \Psi')$ and obtain a reduced diagonal complex (Ω'_1, d'_1) which is a diagonal complex with rotation constant $C_\Psi + C_\Phi - R_D$. By proposition 3.7, applying lemmas 3.1, 3.2, and 3.3 do not change the configuration of the right lower block in the matrices of equation (14). Thus, the complex $D(\Phi, \Psi)$ is homotopy equivalent to a complex with segments

$$(15) \quad \dots \begin{bmatrix} \Omega_1^{r-1} \\ \Omega_n^{s-1} \end{bmatrix} \xrightarrow{\begin{pmatrix} d_1^{r-1} & 0 \\ \rho_1^{r-1} & d_n^{s-1} \end{pmatrix}} \begin{bmatrix} \Omega_1^r \\ \Omega_n^s \end{bmatrix} \xrightarrow{\begin{pmatrix} d_1^r & 0 \\ \rho_1^r & d_n^s \end{pmatrix}} \begin{bmatrix} \Omega_1^{r+1} \\ \Omega_n^{s+1} \end{bmatrix} \xrightarrow{\begin{pmatrix} d_1^{r+1} & 0 \\ \rho_1^{r+1} & d_n^{s+1} \end{pmatrix}} \begin{bmatrix} \Omega_1^{r+2} \\ \Omega_n^{s+2} \end{bmatrix} \dots$$

that only have loops in the column blocks Ω_n^s .

According to proposition 5.8 the chain complex $D(\Psi, \phi_N) = \dots \Omega_n^{s-1} \xrightarrow{d_n^{s-1}} \Omega_n^s \xrightarrow{d_n^s} \Omega_n^{s+1} \dots$ is a coherently diagonal complex with rotation constant $C_\Psi - \overline{R}(\phi_N) - R_D$, then for each homological degree s of (Ψ, e) , and each smoothing ψ in Ψ^s , we have that $2s - R(D(\psi, \phi_N)) = C_\Psi - R(\phi_N) - R_D$. Adding $2t_M$ to each of these last equations we obtain $2r - R(D(\psi, \phi_N)) = C_\Psi + C_\Phi - R_D$. That proves that we can obtain a reduced diagonal complex from (Ω, d) . \square

Proof. (Of **Theorem 1**) By proposition 4.6 we just need to prove that \mathcal{D} is closed under composition of basic operators. Let $(\Omega_1, d_1) \in \mathcal{D}$ and let U be a basic unary operator. Since $U(\Omega_1)$ is a partial closure of (Ω_1, d_1) , $U(\Omega_1)$ is diagonal. Furthermore any partial closure of $U(\Omega_1)$ is also a partial closure of (Ω_1, d_1) , so $U(\Omega_1) \in \mathcal{D}$.

Let (Ω_1, d_1) and (Ω_2, d_2) be elements of \mathcal{D} and D a basic binary operator. Since D is defined from a type- \mathcal{A} diagram, there is at least one boundary arc. Without loss of generality, we can assume that there is one boundary arc ending in the first input disc of D . By proposition 5.6, $D(\Omega_1, \Omega_2)$ is alternating. Let $C(D(\Omega_1, \Omega_2))$ be a partial closure of $D(\Omega_1, \Omega_2)$, by proposition 5.6 there exist $P', Q' \in \mathcal{D}$ and a binary operator D' defined from a no-curl planar diagram such that $C(D(\Omega_1, \Omega_2)) = D'(\Omega'_1, \Omega'_2)$. By using now Lemma 6.1 we obtain that $D'(\Omega'_1, \Omega'_2)$ is a diagonal complex. \square

7. NON-SPLIT ALTERNATING TANGLES AND LEE'S THEOREM

7.1. Proof of Theorem 2. We use the concept of gravity information introduced in [Bur] to give an alternating orientation in the boundaries of the tangles.

Proof. (Of **Theorem 2**) The Khovanov complex of a 1-crossing tangle is an element of $\mathcal{D}(2)$. See first example 5.3. Any non-split alternating k -strand tangle with n crossing T , is obtained by a composition of n of these 1-crossing tangles, T_1, \dots, T_n , in a n -input type- \mathcal{A} planar diagram. Any non-split alternating k -strand tangle T with n crossing, is obtained by a composition of n of these 1-crossing tangles, T_1, \dots, T_n , in an n -input type- \mathcal{A} planar diagram. Since the Khovanov homology is a planar algebra morphism, using the same n -input planar diagram for composing $Kh(T_1), \dots, Kh(T_n)$ we obtain the Khovanov homology of the original tangle. By Theorem 1, this is a complex in \mathcal{D} . \square

Corollary 7.1. *The Khovanov complex $[T]$ of a non-split alternating 1-tangle T is homotopy equivalent to a complex*

$$\dots \longrightarrow \Omega^r \{2r + K\} \longrightarrow \Omega^{r+1} \{2(r+1) + K\} \longrightarrow \dots$$

where every Ω^r is a vector of single lines, and K is a constant.

Proof. We just have to apply theorem 2 and see that the rotation number of a 1 open arc, which is the only simple possible smoothing resulting from a 1-tangle, is zero. \square

Figure 8 shows a diagonal complex which is obtained from embedding the complex in Figure 7 in a positive unary basic planar diagram, and then applying lemmas 3.1 and 3.2. The smoothings in this complex have only one strand. Since the rotation number of a smoothing with a unique strand is always 0, we have that the degree shift and the homological degree multiplied by two are in a single diagonal, i.e., $2r - q_r$ is a constant.




Figure 8. A diagonal complex with only one strand in each of its smoothings.

Corollary 7.2. (*Lee's theorem*) *The Khovanov complex $Kh(L)$ of a non-split alternating Link L is homotopy equivalent to a complex:*

$$\cdots \longrightarrow \begin{pmatrix} \Phi^r \{q_r + 1\} \\ \Phi^r \{q_r - 1\} \end{pmatrix} \longrightarrow \begin{pmatrix} \Phi^{r+1} \{q_{r+1} + 1\} \\ \Phi^{r+1} \{q_{r+1} - 1\} \end{pmatrix} \longrightarrow \cdots$$

where every Φ^r is a matrix of empty 1-manifolds, $q_r = 2r + K$, K a constant, and every differential is a matrix in \mathbb{Q} .

Proof. Every non-split alternating link L is obtained by putting a 1-strand tangle T in a 1-input planar diagram with no boundary. Hence, by applying the operator defined from this 1-input planar diagrams to the Khovanov complex of this 1-strand tangle we obtain the Khovanov complex of a link L . By doing that, the vectors of open arcs that we have in corollary 7.1 become vectors of circles. Moreover, every cobordism of the complex transforms in a multiple of a dotted cylinder. Thus, using Lemma 3.1 converts every single loop in a pair of empty sets $\emptyset\{r + K + 1\}$, $\emptyset\{r + K - 1\}$ and every dotted cylinder in an element of the ground field. \square

Figure 9 displays the closure of the complex in Figure 8. After applying lemmas 3.1 and 3.2 we obtain the complex supported in two lines displayed on Figure 9. That is what Lee's Theorem states.

Remark 7.3. It is clear that if our ground ring is \mathbb{Q} , as in the case of [Lee], we can use repeatedly lemma 3.2 in the complex in corollary 7.2 and obtain from this complex, one whose differentials are zero, i.e, the Khovanov homology of the link.




Figure 9. A closure of a coherently diagonal complex is a width-two complex.

REFERENCES

- [BN] D. BAR-NATAN, *On Khovanov's categorification of de Jones polynomial*, Alg. Geom. Top., **2** (2002), 337-370.
- [BN1] D. BAR-NATAN, *Khovanov's homology for tangles and cobordisms*, Geometry & Topology, **9** (2005), 1443-1499.
- [BN2] D. BAR-NATAN, *Fast Khovanov homology computations*, arXiv:math.GT/0606318 (2006)
- [BN3] D. BAR-NATAN AND S. MORRISON, *The Karoubi envelope and Lee's degeneration of Khovanov homology*, Algebraic & Geometry Topology, **6** (2006), 1459-1469.
- [Bur] H. BURGOS SOTO, *The Jones polynomial and the planar algebra of alternating links*, arXiv:math.GT/0807.2600v1 (2008)
- [Ga] S. GAROUFALIDIS, *A conjecture on Khovanov's invariants*, Fundamenta Mathematicae, **184** (2001), 99-101.
- [Jo] V. JONES, *Planar Algebras, I*, New Zealand Journal of Mathematics., **bbb** (200), aa-bb.
- [Kh] M. KHOVANOV, *A categorification of the Jones Polynomial*, Duke Math J., **101(3)** (1999), 359-426.
- [Lee] E. S. LEE, *The support of the Khovanov's invariants for alternating links*, arXiv:math.GT/0201105, v1, (2002).
- [Lee1] E. S. LEE, *A Khovanov invariant for alternating links*, arXiv:math.GT/0210213, v2, (2002).
- [Naot] G. NAOT, *On the Algebraic Structure of Bar-Natan's Universal Complex and the Geometric Structure of Khovanov Link Homology Theories*, arXiv:math.GT/0603347 (2006)
- [Thi] M. THISTLETHWAITE, *Spanning tree expansion of the Jones polynomial*, Topology, **26** (1987), 297-309.
- [Wei] C. A. WEIBEL, *An Introduction to Homological Algebra*, Cambridge Studies in Advanced Mathematics 38, Cambridge University Press, Cambridge, UK., (1994).

SCHOOL OF ENGINEERING TECHNOLOGY & APPLIED SCIENCE, CENTENNIAL COLLEGE, TORONTO, ON, M1C 5J9, CANADA

E-mail address: hburgos1@my.centennialcollege.ca

URL: <http://individual.utoronto.ca/hernandoburgos/>